

Roberto, K.A., Perkins, S.N., & Holland, A. K. (2007)  
Research on Persistent Pain in Late Life: Current Topics and Challenges  
*Journal of Women & Aging, 19* (3/4)

*Articles Reviewed*

- Altman, R. D., Akermark, C., Beaulieu, A. D., & Schnitzer, T. (2004). Efficacy and safety of a single intra-articular injection of non-animal stabilized hyaluronic acid (NASHA) in patients with osteoarthritis of the knee. *Osteoarthritis Cartilage, 12*(8), 642-649.
- Amin, S., Luepongsak, N., McGibbon, C. A., LaValley, M. P., Krebs, D. E., & Felson, D. T. (2004). Knee adduction moment and development of chronic knee pain in elders. *Arthritis and Rheumatism, 51*(3), 371-376.
- Averbuch, M., & Katzper, M. (2004). Assessment of visual analog versus categorical scale for measurement of osteoarthritis pain. *Journal of Clinical Pharmacology, 44*(4), 368-372.
- Baier, R. R., Gifford, D. R., Patry, G., Banks, S. M., Rochon, T., DeSilva, D., et al. (2004). Ameliorating pain in nursing homes: A collaborative quality-improvement project. *Journal of the American Geriatric Society, 52*(12), 1988-1995.
- Bailey, R. L., Ledikwe, J. H., Smiciklas-Wright, H., Mitchell, D. C., & Jensen, G. L. (2004). Persistent oral health problems associated with comorbidity and impaired diet quality in older adults. *Journal of the American Dietetic Association, 104*(8), 1273-1276.
- Baird, C. L. (2000). Living with hurting and difficulty doing: older women with osteoarthritis. *Clinical Excellence for Nurse Practitioners: The International Journal of NPACE, 4*(4), 231-237.
- Baird, C. L., & Sands, L. (2004). A pilot study of the effectiveness of guided imagery with progressive muscle relaxation to reduce chronic pain and mobility difficulties of osteoarthritis. *Pain Management Nursing, 5*(3), 97-104.
- Baker, T. A. (2003). Arthritis symptoms as indicators of pain in older African Americans. *Ethnicity & Disease, 13*(4), 513-520.
- Barry, L. C., Guo, Z., Kerns, R. D., Duong, B. D., & Reid, M. C. (2003). Functional self-efficacy and pain-related disability among older veterans with chronic pain in a primary care setting. *Pain, 104*(1-2), 131-137.
- Barry, L. C., Kerns, R. D., Guo, Z., Duong, B. D., Iannone, L. P., & Carrington Reid, M. (2004). Identification of strategies used to cope with chronic pain in older persons receiving primary care from a Veterans Affairs Medical Center. *Journal of the American Geriatric Society, 52*(6), 950-956.
- Battisti, E., Piazza, E., Rigato, M., Nuti, R., Bianciardi, L., Scribano, A., & Giordano, N. (2004). Efficacy and safety of a musically modulated electromagnetic field (TAMMEF) in patients affected by knee osteoarthritis. *Clinical and Experimental Rheumatology, 22*(5), 568-572.

- Bennett, J. A., Stewart, A. L., Kayser-Jones, J., & Glaser, D. (2002). The mediating effect of pain and fatigue on level of functioning in older adults. *Nursing Research, 51*(4), 254-265.
- Berger, A., Dukes, E. M., & Oster, G. (2004). Clinical characteristics and economic costs of patients with painful neuropathic disorders. *Journal of Pain, 5*(3), 143-149.
- Bergeron, S., Ommen, S. R., Bailey, K. R., Oh, J. K., McCully, R. B., & Pellikka, P. A. (2004). Exercise echocardiographic findings and outcome of patients referred for evaluation of dyspnea. *Journal of the American College of Cardiology, 43*(12), 2242-2246.
- Berman, B. M., Lao, L., Langenberg, P., Lee, W. L., Gilpin, A. M., & Hochberg, M. C. (2004). Effectiveness of acupuncture as adjunctive therapy in osteoarthritis of the knee: A randomized, controlled trial. *Annals of Internal Medicine, 141*(12), 901-910.
- Berry, J. D., Rowbotham, M. C., & Petersen, K. L. (2004). Complex regional pain syndrome-like symptoms during herpes zoster. *Pain, 110*(1-2), 8-9.
- Bishop, K. L., Ferraro, F., & Borowiak, D. M. (2001). Pain management in older adults: Role of fear and avoidance. *Clinical Gerontologist, 23*(1-2), 33-42.
- Blixen, C. E., & Kippes, C. (1999). Depression, social support, and quality of life in older adults with osteoarthritis. *Image: The Journal of Nursing Scholarship, 31*(3), 221-226.
- Bookwala, J., Harralson, T. L., & Parmelee, P. A. (2003). Effects of pain on functioning and well-being in older adults with osteoarthritis of the knee. *Psychology and Aging, 18*(4), 844-850.
- Bosley, B. N., Weiner, D. K., Rudy, T. E., & Granieri, E. (2004). Is chronic nonmalignant pain associated with decreased appetite in older adults? Preliminary evidence. *Journal of the American Geriatric Society, 52*(2), 247-251.
- Brandt, K. D., Heilman, D. K., Slemenda, C., Katz, B. P., Mazzuca, S., Braunstein, E. M., & Byrd, D. (2000). A comparison of lower extremity muscle strength, obesity, and depression scores in elderly subjects with knee pain with and without radiographic evidence of knee osteoarthritis. *The Journal of Rheumatology, 27*(8), 1937-1946.
- Bruce, B., & Fries, J. (2004). Longitudinal comparison of the Health Assessment Questionnaire (HAQ) and the Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC). *Arthritis and Rheumatism, 51*(5), 730-737.
- Brummel-Smith, K., London, M. R., Drew, N., Krulewitch, H., Singer, C., & Hanson, L. (2002). Outcomes of pain in frail older adults with dementia. *Journal of the American Geriatrics Society, 50*(11), 1847-1851.
- Caborn, D., Rush, J., Lanzer, W., Parenti, D., & Murray, C. (2004). A randomized, single-blind comparison of the efficacy and tolerability of hylan G-F 20 and triamcinolone hexacetonide in patients with osteoarthritis of the knee. *The Journal of Rheumatology, 31*(2), 333-343.

- Cadogan, M. P., Schnelle, J. F., Yamamoto-Mitani, N., Cabrera, G., & Simmons, S. F. (2004). A minimum data set prevalence of pain quality indicator: is it accurate and does it reflect differences in care processes? *The Journal of Gerontology: Medical Sciences*, 59(3), M281-M285.
- Carbone, L. D., Nevitt, M. C., Wildy, K., Barrow, K. D., Harris, F., Felson, D., Peterfy, C., Visser, M., Harris, T. B., Wang, B. W. E., & Kritchevsky, S. B. (2004). The relationship of antiresorptive drug use to structural findings and symptoms of knee osteoarthritis. *Arthritis and Rheumatism*, 50(11), 3516-3525.
- Carels, R. A., Musher-Eizenman, D., Cacciapaglia, H., Perez-Benitez, C. I., Christie, S., & O'Brien, W. (2004). Psychosocial functioning and physical symptoms in heart failure patients: A within-individual approach. *Journal of Psychosomatic Research*, 56(1), 95-101.
- Carr, M. P., France, R. P., & Horton, J. E. (2002). A comparison of perceived injection pain with and without anesthetic between elderly and young female subjects. *Special Care in Dentistry*, 22(2), 75-79.
- Carrington Reid, M., Williams, C. S., Concato, J., Tinetti, M. E., & Gill, T. M. (2003). Depressive symptoms as a risk factor for disabling back pain in community-dwelling older persons. *Journal of the American Geriatric Society*, 51(12), 1710-1717.
- Carrington Reid, M., Williams, C. S., & Gill, T. M. (2003). The relationship between psychological factors and disabling musculoskeletal pain in community-dwelling older persons. *Journal of the American Geriatric Society*, 51(8), 1092-1098.
- Casarett, D. J., Hirschman, K. B., Miller, E. R., & Farrar, J. T. (2002). Is satisfaction with pain management a valid and reliable quality indicator for use in nursing homes? *Journal of the American Geriatric Society*, 50(12), 2029-2034.
- Chandra, H. R., Goldstein, J. A., Choudhary, N., O'Neill, C. S., George, P. B., Gangasani, S. R., Cronin, L., Marcovitz, P. A., Hauser, A. M., & O'Neill, W. W. (2004). Adverse outcome in aortic sclerosis is associated with coronary artery disease and inflammation. *Journal of the American College of Cardiology*, 43(2), 169-175.
- Chang, A., Hayes, K., Dunlop, D., Hurwitz, D., Song, J., Cahue, S., Genge, R., & Sharma, L. (2004). Thrust during ambulation and the progression of knee osteoarthritis. *Arthritis and Rheumatism*, 50(12), 3897-3903.
- Chibnall, J. T., & Tait, R. C. (2001). Pain assessment in cognitively impaired and unimpaired older adults: A comparison of four scales. *Pain*, 92(1-2), 173-186.
- Chodosh, J., Ferrell, B. A., Shekelle, P. G., & Wenger, N. S. (2001). Quality indicators for pain management in vulnerable elders. *Annals of Internal Medicine*, 135, 731-735.
- Chodosh, J., Solomon, D. H., Roth, C. P., Chang, J. T., MacLean, C. H., Ferrell, B. A., Shekelle, P. G., & Wenger, N. S. (2004). The quality of medical care provided to vulnerable older patients with chronic pain. *Journal of the American Geriatric Society*, 52(5), 756-761.

- Christmas, C., Crespo, C. J., Franckowiak, S. C., Bathon, J. M., Bartlett, S. J., & Andersen, R. E. (2002). How common is hip pain among older adults? Results from the Third National Health and Nutrition Examination Survey. *Journal of Family Practice*, 51(4), 345-348.
- Chu, L., Schnelle, J. F., Cadogan, M. P., & Simmons, S. F. (2004). Using the minimum data set to select nursing home residents for interview about pain. *Journal of the American Geriatric Society*, 52(12), 2057-2061.
- Cibere, J., Kopec, J. A., Thorne, A., Singer, J., Canvin, J., Robinson, D. B., Pope, J., Hong, P., Grant, E., & Esdaile, J. M. (2004). Randomized, double-blind, placebo-controlled glucosamine discontinuation trial in knee osteoarthritis. *Arthritis and Rheumatism*, 51(5), 738-745.
- Clark, L., Jones, K., & Pennington, K. (2004). Pain assessment practices with nursing home residents. *Western Journal of Nursing Research*, 26(7), 733-750.
- Cohen-Mansfield, J. (2002). Relatives' assessment of pain in cognitively impaired nursing home residents. *Journal of Pain and Symptom Management*, 24(6), 562-571.
- Cohen-Mansfield, J. (2004). The adequacy of the minimum data set assessment of pain in cognitively impaired nursing home residents. *Journal of Pain and Symptom Management*, 27(4), 343-351.
- Cohen-Mansfield, J., & Lipson, S. (2002). Pain in cognitively impaired nursing home residents: How well are physicians diagnosing it? *Journal of the American Geriatrics Society*, 50(6), 1039-1044.
- Cook, A. J., & Chastain, D. C. (2001). The classification of patients with chronic pain: Age and sex differences. *Pain Research & Management : The Journal of the Canadian Pain Society*, 6(3), 142-151.
- Coplan, P. M., Schmader, K., Nikas, A., Chan, I. S. F., Choo, P., Levin, M. J., Johnson, G., Bauer, M., Williams, H. M., Kaplan, K. M., Guess, H. A., & Oxman, M. N. (2004). Development of a measure of the burden of pain due to herpes zoster and postherpetic neuralgia for prevention trials: Adaptation of the Brief Pain Inventory. *Journal of Pain*, 5(6), 344-356.
- Coppola, K. M., Bookwala, J., Ditto, P. H., Lockhart, L. K., Danks, J. H., & Smucker, W. D. (1999). Elderly adults' preferences for life-sustaining treatments: The role of impairment, prognosis, and pain. *Death Studies*, 23(7), 617-634.
- Creamer, P., Lethbridge-Cejku, M., Costa, P., Tobin, J. D., Herbst, J. H., & Hochberg, M. C. (1999). The relationship of anxiety and depression with self-reported knee pain in the community: Data from the Baltimore Longitudinal Study of Aging. *Arthritis Care and Research*, 12(1), 3-7.
- Cremeans-Smith, J. K., Stephens, M. A. P., Franks, M. M., Martire, L. M., Druley, J. A., & Wojno, W. C. (2003). Spouses' and physicians' perceptions of pain severity in older women with osteoarthritis: Dyadic agreement and patients' well-being. *Pain*, 106(1-2), 27-34.

- Currie, S. R., & Wang, J. (2004). Chronic back pain and major depression in the general Canadian population. *Pain, 107*(1-2), 54-60.
- Davis, G. C., Hiemenz, M. L., & White, T. L. (2002). Barriers to managing chronic pain of older adults with arthritis. *Journal of Nursing Scholarship, 34*(2), 121-126.
- Day, C. S., Gelberman, R., Patel, A. A., Vogt, M. T., Ditsios, K., & Boyer, M. I. (2004). Basal joint osteoarthritis of the thumb: A prospective trial of steroid injection and splinting. *Journal of Hand Surgery, 29*(2), 247-251.
- DeVon, H. A., & Zerwic, J. J. (2004). Differences in the symptoms associated with unstable angina and myocardial infarction. *Progress in Cardiovascular Nursing, 19*(1), 6-11.
- Dominick, K. L., Bosworth, H. B., Dudley, T. K., Waters, S. J., Campbell, L. C., & Keefe, F. J. (2004). Patterns of opioid analgesic prescription among patients with osteoarthritis. *Journal of Pain & Palliative Care Pharmacotherapy, 18*(1), 31-46.
- Dominick, K. L., Bosworth, H. B., Hsieh, J. B., & Moser, B. K. (2004). Racial differences in analgesic/anti-inflammatory medication use and perceptions of efficacy. *Journal of the National Medical Association, 96*(7), 928-932.
- Dominick, K. L., Bosworth, H. B., Jeffreys, A. S., Grambow, S. C., Oddone, E. Z., & Horner, R. D. (2004). Racial/ethnic variations in non-steroidal anti-inflammatory drug (NSAID) use among patients with osteoarthritis. *Pharmacoepidemiology and Drug Safety, 13*(10), 683-694.
- Druley, J. A., Stephens, M. A., Martire, L. M., Ennis, N., & Wojno, W. C. (2003). Emotional congruence in older couples coping with wives' osteoarthritis: Exacerbating effects of pain behavior. *Psychology and Aging, 18*(3), 406-414.
- Dunn, K. S., & Horgas, A. L. (2004). Religious and nonreligious coping in older adults experiencing chronic pain. *Pain Management Nursing, 5*(1), 19-28.
- Emkey, R., Rosenthal, N., Wu, S. C., Jordan, D., & Kamin, M. (2004). Efficacy and safety of tramadol/acetaminophen tablets (Ultracet) as add-on therapy for osteoarthritis pain in subjects receiving a COX-2 nonsteroidal antiinflammatory drug: a multicenter, randomized, double-blind, placebo-controlled trial. *The Journal of Rheumatology, 31*(1), 150-156.
- Engle, V. F., Graney, M. J., & Chan, A. (2001). Accuracy and bias of licensed practical nurse and nursing assistant ratings of nursing home residents' pain. *The Journal of Gerontology: Medical Sciences, 56*(7), M405-M411.
- Ersek, M., Turner, J. A., Cain, K. C., & Kemp, C. A. (2004). Chronic pain self-management for older adults: A randomized controlled trial. *BMC Geriatrics, 4*(1), 4-7.
- Ersek, M., Turner, J. A., McCurry, S. M., Gibbons, L., & Kraybill, B. M. (2003). Efficacy of a self-management group intervention for elderly persons with chronic pain. *The Clinical Journal of Pain, 19*(3), 156-167.

- Ferrell, B. A., Stein, W. M., & Beck, J. C. (2000). The Geriatric Pain Measure: Validity, reliability and factor analysis. *Journal of the American Geriatrics Society*, 48(12), 1669-1673.
- Fisher, S. E., Burgio, L. D., Thorn, B. E., Allen-Burge, R., Gerstle, J., Roth, D. L., & Allen, S. J. (2002). Pain assessment and management in cognitively impaired nursing home residents: Association of certified nursing assistant pain report, Minimum Data Set pain report, and analgesic medication use. *Journal of the American Geriatrics Society*, 50(1), 152-156.
- Fortess, E. E., Soumerai, S. B., McLaughlin, T. J., & Ross-Degnan, D. (2001). Utilization of essential medications by vulnerable older people after a drug benefit cap: importance of mental disorders, chronic pain, and practice setting. *Journal of the American Geriatrics Society*, 49(6), 793-797.
- Gagliese, L., & Melzack, R. (2003). Age-related differences in the qualities but not the intensity of chronic pain. *Pain*, 104(3), 597-608.
- Gaines, J. M., Metter, E. J., & Talbot, L. A. (2004). The effect of neuromuscular electrical stimulation on arthritis knee pain in older adults with osteoarthritis of the knee. *Applied Nursing Research*, 17(3), 201-206.
- Gardner, A. W., Montgomery, P. S., & Killewich, L. A. (2004). Natural history of physical function in older men with intermittent claudication. *Journal of Vascular Surgery*, 40(1), 73-78.
- Gibson, S. J., & Helme, R. D. (2000). Cognitive factors and the experience of pain and suffering in older persons. *Pain*, 85(3), 375-383.
- Gloth, F. M., 3rd, Scheve, A. A., Stober, C. V., Chow, S., & Prosser, J. (2001). The Functional Pain Scale: Reliability, validity, and responsiveness in an elderly population. *Journal of the American Medical Directors Association*, 2(3), 110-114.
- Graves, N., Krepcho, M., Mayo, H. G., & Hill, J. (2004). Clinical inquiries. Does yoga speed healing for patients with low back pain? *Journal of Family Practice*, 53(8), 661-662.
- Green, C. R., Baker, T. A., Smith, E. M., & Sato, Y. (2003). The effect of race in older adults presenting for chronic pain management: A comparative study of black and white Americans. *Journal of Pain*, 4(2), 82-90.
- Hadjistavropoulos, T., LaChapelle, D. L., MacLeod, F. K., Snider, B., & Craig, K. D. (2000). Measuring movement-exacerbated pain in cognitively impaired frail elders. *The Clinical Journal of Pain*, 16(1), 54-63.
- Harrison, A. L. (2004). The influence of pathology, pain, balance, and self-efficacy on function in women with osteoarthritis of the knee. *Physical Therapy*, 84(9), 822-831.
- Hasegawa, H. (2004). Utilization of zonisamide in patients with chronic pain or epilepsy refractory to other treatments: A retrospective, open label, uncontrolled study in a VA hospital. *Current Medical Research and Opinion*, 20(5), 577-580.

- Hawk, C., Long, C. R., Boulanger, K. T., Morschhauser, E., & Fuhr, A. W. (2000). Chiropractic care for patients aged 55 years and older: Report from a practice-based research program. *Journal of the American Geriatrics Society, 48*(5), 534-545.
- Hirsch, R., Guralnik, J. M., Leveille, S. G., Simonsick, E. M., Ling, S., Bandeen-Roche, K., Rantanen, T., Pahor, M., Fried, L. P., & Hochberg, M. C. (1999). Severity of hand osteoarthritis and its association with upper extremity impairment in a population of disabled older women: The Women's Health and Aging Study. *Aging, 11*(4), 253-261.
- Horgas, A. L., & Dunn, K. (2001). Pain in nursing home residents. Comparison of residents' self-report and nursing assistants' perceptions. Incongruencies exist in resident and caregiver reports of pain; therefore, pain management education is needed to prevent suffering. *Journal of Gerontological Nursing, 27*(3), 44-53.
- Horowitz, B. P., & Chang, P. F. J. (2004). Promoting well-being and engagement in life through occupational therapy lifestyle redesign: A pilot study within adult day programs. *Topics in Geriatric Rehabilitation, 20*(1), 46-58.
- Hughes, S. L., Seymour, R. B., Campbell, R., Pollak, N., Huber, G., & Sharma, L. (2004). Impact of the fit and strong intervention on older adults with osteoarthritis. *The Gerontologist, 44*(2), 217-228.
- Ibrahim, S. A., Burant, C. J., Mercer, M. B., Siminoff, L. A., & Kwok, C. K. (2003). Older patients' perceptions of quality of chronic knee or hip pain: Differences by ethnicity and relationship to clinical variables. *The Journal of Gerontology: Medical Sciences, 58*(5), M472-M477.
- Inadequate Pain Care for Elders: The Need for a Primary Care-Pain Medicine Community Collaboration. (2002). *Pain Medicine, 3*(2), 180.
- Iversen, M. D., Fossel, A. H., & Katz, J. N. (2003). Enhancing function in older adults with chronic low back pain: A pilot study of endurance training. *Archives of Physical Medicine and Rehabilitation, 84*(9), 1324-1331.
- Jenq, G. Y., Guo, Z., Drickamer, M., Marottoli, R. A., & Reid, M. C. (2004). Timing in the communication of pain among nursing home residents, nursing staff, and clinicians. *Archives of Internal Medicine, 164*(14), 1508-1512.
- Kaasalainen, S., & Crook, J. (2003). A comparison of pain-assessment tools for use with elderly long-term-care residents. *Canadian Journal of Nursing Research, 35*(4), 58-71.
- Kaasalainen, S., & Crook, J. (2004). An exploration of seniors' ability to report pain. *Clinical Nursing Research, 13*(3), 199-215.
- Kamel, H. K., Phlavan, M., Malekgoudarzi, B., Gogel, P., & Morley, J. E. (2001). Utilizing pain assessment scales increases the frequency of diagnosing pain among elderly nursing home residents. *Journal of Pain Symptom Manage, 21*(6), 450-455.
- Kane, R. L., Homyak, P., Bershady, B., Lum, Y. S., & Siadat, M. S. (2003). Outcomes of managed care of dually eligible older persons. *The Gerontologist, 43*(2), 165-174.

- Kaplan, M. S., Huguette, N., Newsom, J. T., & McFarland, B. H. (2003). Characteristics of physically inactive older adults with arthritis: Results of a population-based study. *Preventive Medicine, 37*(1), 61-67.
- Keefe, F. J., Affleck, G., France, C. R., Emery, C. F., Waters, S., Caldwell, D. S., Stainbrook, D., Hackshaw, K. V., Fox, L. C., & Wilson, K. (2004). Gender differences in pain, coping, and mood in individuals having osteoarthritic knee pain: A within-day analysis. *Pain, 110*(3), 571-577.
- Keefe, F. J., Affleck, G., Lefebvre, J., Underwood, L., Caldwell, D. S., Drew, J., Egert, J., Gibson, J., & Pargament, K. (2001). Living with rheumatoid arthritis: The role of daily spirituality and daily religious and spiritual coping. *Journal of Pain, 2*(2), 101-110.
- Keefe, F. J., Blumenthal, J., Baucom, D., Affleck, G., Waugh, R., Caldwell, D. S., Beaupre, P., Kashikar-Zuck, S., Wright, K., Egert, J., & Lefebvre, J. (2004). Effects of spouse-assisted coping skills training and exercise training in patients with osteoarthritic knee pain: A randomized controlled study. *Pain, 110*(3), 539-549.
- Kenefick, A. L. (2004). Pain treatment and quality of life: Reducing depression and improving cognitive impairment. *Journal of Gerontological Nursing, 30*(5), 22-29.
- Klinger, L., Spaulding, S. J., Polatajko, H. J., MacKinnon, J. R., & Miller, L. (1999). Chronic pain in the elderly: occupational adaptation as a means of coping with osteoarthritis of the hip and/or knee. *Clinical Journal of Pain, 15*(4), 275-283.
- Krusewitz, H., London, M. R., Skakel, V. J., Lundstedt, G. J., Thomason, H., & Brummel-Smith, K. (2000). Assessment of pain in cognitively impaired older adults: A comparison of pain assessment tools and their use by nonprofessional caregivers. *Journal of the American Geriatrics Society, 48*(12), 1607-1611.
- Kunstler, R., Greenblatt, F., & Moreno, N. (2004). Aromatherapy and hand massage: Therapeutic recreation interventions for pain management. *Therapeutic Recreation Journal, 38*(2), 133-147.
- Lamb, S. E., Guralnik, J. M., Buchner, D. M., Ferrucci, L. M., Hochberg, M. C., Simonsick, E. M., & Fried, L. P. (2000). Factors that modify the association between knee pain and mobility limitation in older women: The Women's Health and Aging Study. *Annals of the Rheumatic Diseases, 59*(5), 331-337.
- Lane, N. E., Nevitt, M. C., Hochberg, M. C., Hung, Y. Y., & Palermo, L. (2004). Progression of radiographic hip osteoarthritis over eight years in a community sample of elderly white women. *Arthritis and Rheumatism, 50*(5), 1477-1486.
- Letizia, M., Creech, S., Norton, E., Shanahan, M., & Hedges, L. (2004). Barriers to caregiver administration of pain medication in hospice care. *Journal of Pain Symptom Manage, 27*(2), 114-124.
- Leveille, S. G., Bean, J., Bandeen-Roche, K., Jones, R., Hochberg, M., & Guralnik, J. M. (2002). Musculoskeletal pain and risk for falls in older disabled women living in the community. *Journal of the American Geriatrics Society, 50*(4), 671-678.



- Leveille, S. G., Fried, L., & Guralnik, J. M. (2002). Disabling symptoms: What do older women report? *Journal of General Internal Medicine*, 17(10), 766-773.
- Leveille, S. G., Fried, L. P., McMullen, W., & Guralnik, J. M. (2004). Advancing the taxonomy of disability in older adults. *The Journal of Gerontology: Medical Sciences*, 59(1), M86-M93.
- Leveille, S. G., Ling, S., Hochberg, M. C., Resnick, H. E., Bandeen-Roche, K. J., Won, A., & Guralnik, J. M. (2001). Widespread musculoskeletal pain and the progression of disability in older disabled women. *Annals of Internal Medicine*, 135(12), 1038-1046.
- Ling, S. M., Fried, L. P., Garrett, E. S., Fan, M. Y., Rantanen, T., & Bathon, J. M. (2003). Knee osteoarthritis compromises early mobility function: The Women's Health and Aging Study II. *The Journal of Rheumatology*, 30(1), 114-120.
- Loeslie, V. (2000). Pain in the elderly: Polymyalgia rheumatica. *Clinical Excellence for Nurse Practitioners : The International Journal of NPACE*, 4(6), 345-348.
- Mangione, K. K., McCully, K., Gloviak, A., Lefebvre, I., Hofmann, M., & Craik, R. (1999). The effects of high-intensity and low-intensity cycle ergometry in older adults with knee osteoarthritis. *The Journal of Gerontology: Medical Sciences*, 54(4), M184-M190.
- Manz, B. D., Mosier, R., Nusser-Gerlach, M. A., Bergstrom, N., & Agrawal, S. (2000). Pain assessment in the cognitively impaired and unimpaired elderly. *Pain Management Nursing*, 1(4), 106-115.
- Marsh, A. P., Rejeski, W. J., Lang, W., Miller, M. E., & Messier, S. P. (2003). Baseline balance and functional decline in older adults with knee pain: The Observational Arthritis Study in Seniors. *Journal of the American Geriatric Society*, 51(3), 331-339.
- Martin, K. A., Rejeski, W. J., Miller, M. E., James, M. K., Ettinger, W. H., Jr., & Messier, S. P. (1999). Validation of the PASE in older adults with knee pain and physical disability. *Medicine and Science in Sports and Exercise*, 31(5), 627-633.
- Mazucca, S. A., Page, M. C., Meldrum, R. D., Brandt, K. D., & Petty-Saphon, S. (2004). Pilot study of the effects of a heat-retaining knee sleeve on joint pain, stiffness, and function in patients with knee osteoarthritis. *Arthritis and Rheumatism*, 51(5), 716-721.
- McAlindon, T., Formica, M., LaValley, M., Lehmer, M., & Kabbara, K. (2004). Effectiveness of glucosamine for symptoms of knee osteoarthritis: Results from an internet-based randomized double-blind controlled trial. *American Journal of Medicine*, 117(9), 643-649.
- McCaffrey, R., & Freeman, E. (2003). Effect of music on chronic osteoarthritis pain in older people. *Journal of Advanced Nursing*, 44(5), 517-524.
- McDermott, M. M., Liu, K., Greenland, P., Guralnik, J. M., Criqui, M. H., Chan, C., Pearce, W. H., Schneider, J. R., Ferrucci, L., Celic, L., Taylor, L. M., Vonesh, E., Martin, G. J., & Clark, E. (2004). Functional decline in peripheral arterial disease: associations with the ankle brachial index and leg symptoms. *Journal of the American Medical Association*, 292(4), 453-461.

- McKiernan, F., Faciszewski, T., & Jensen, R. (2004). Quality of life following vertebroplasty. *Journal of Bone and Joint Surgery*, 86-A(12), 2600-2606.
- Menard, J. R., Smith, H. E., Riebe, D., Braun, C. M., Blissmer, B., & Patterson, R. B. (2004). Long-term results of peripheral arterial disease rehabilitation. *Journal of Vascular Surgery*, 39(6), 1186-1192.
- Mentes, J. C., Teer, J., & Cadogan, M. P. (2004). The pain experience of cognitively impaired nursing home residents: Perceptions of family members and certified nursing assistants. *Pain Management Nursing*, 5(3), 118-125.
- Messier, S. P., Glasser, J. L., Ettinger, W. H., Jr., Craven, T. E., & Miller, M. E. (2002). Declines in strength and balance in older adults with chronic knee pain: A 30-month longitudinal, observational study. *Arthritis and Rheumatism*, 47(2), 141-148.
- Messier, S. P., Loeser, R. F., Miller, G. D., Morgan, T. M., Rejeski, W. J., Sevick, M. A., Ettinger, W. H., Pahor, M., & Williamson, J. D. (2004). Exercise and dietary weight loss in overweight and obese older adults with knee osteoarthritis: The Arthritis, Diet, and Activity Promotion Trial. *Arthritis and Rheumatism*, 50(5), 1501-1510.
- Middaugh, S. J., & Pawlick, K. (2002). Biofeedback and behavioral treatment of persistent pain in the older adult: A review and a study. *Applied Psychophysiology and Biofeedback*, 27(3), 185-202.
- Miller, M. E., Rejeski, W. J., Messier, S. P., & Loeser, R. F. (2001). Modifiers of change in physical functioning in older adults with knee pain: The Observational Arthritis Study in Seniors (OASIS). *Arthritis and Rheumatism*, 45(4), 331-339.
- Miller, R. R., Zhang, Y., Silliman, R. A., Hayes, M. K., Gleville, S., Murabito, J. M., Kiel, D., O'Connor, G. T., & Felson, D. T. (2004). Effect of medical conditions on improvement in self-reported and observed functional performance of elders. *Journal of the American Geriatric Society*, 52(2), 217-223.
- Mossey, J. M., & Gallagher, R. M. (2004). The longitudinal occurrence and impact of comorbid chronic pain and chronic depression over two years in continuing care retirement community residents. *Pain Medicine*, 5(4), 335-348.
- Mossey, J. M., Gallagher, R. M., & Tirumalasetti, F. (2000). The effects of pain and depression on physical functioning in elderly residents of a continuing care retirement community. *Pain Medicine*, 1(4), 340-350.
- Nair, C. K., Khan, I. A., Mehta, N. J., Ryschon, K. L., & Nair, R. C. (2004). Gender significance of ST-segment deviation detected by ambulatory (Holter) monitoring. *International Journal of Cardiology*, 95(2-3), 153-157.
- Nemeth, K. A., Harrison, M. B., Graham, I. D., & Burke, S. (2004). Understanding venous leg ulcer pain: Results of a longitudinal study. *Ostomy/Wound Management*, 50(1), 34-46.

- Ohldin, A., Young, B., Derleth, A., McDonell, M., Diehr, P., Kiefe, C., & Fihn, S. (2004). Ethnic differences in satisfaction and quality of life in veterans with ischemic heart disease. *Journal of the National Medical Association, 96*(6), 799-808.
- Olson, C. J. (2004). Relief of ischemic leg pain with intermittent positive pressure. *Canadian Family Physician, 50*, 1225-1226.
- Oster, G., Berger, A., Dukes, E., Edelsberg, J., & McCarberg, B. (2004). Use of potentially inappropriate pain-related medications in older adults with painful neuropathic disorders. *The American Journal of Geriatric Pharmacotherapy, 2*(3), 163-170.
- Pahor, M., Guralnik, J. M., Wan, J. Y., Ferrucci, L., Penninx, B. W., Lyles, A., Ling, S., & Fried, L. P. (1999). Lower body osteoarticular pain and dose of analgesic medications in older disabled women: The Women's Health and Aging Study. *American Journal of Public Health, 89*(6), 930-934.
- Proctor, W. R., & Hirdes, J. P. (2001). Pain and cognitive status among nursing home residents in Canada. *Pain Research & Management : The Journal of the Canadian, 6*(3), 119-125.
- Rapp, S. R., Rejeski, W. J., & Miller, M. E. (2000). Physical function among older adults with knee pain: The role of pain coping skills. *Arthritis Care and Research, 13*(5), 270-279.
- Reid, M. C., Guo, Z., Towle, V. R., Kerns, R. D., & Concato, J. (2002). Pain-related disability among older male veterans receiving primary care. *The Journal of Gerontology. Medical Sciences, 57*(11), M727-M732.
- Reid, M. C., Otis, J., Barry, L. C., & Kerns, R. D. (2003). Cognitive-behavioral therapy for chronic low back pain in older persons: A preliminary study. *Pain Medicine, 4*(3), 223-230.
- Reid, M. C., Williams, C. S., & Gill, T. M. (2003). The relationship between psychological factors and disabling musculoskeletal pain in community-dwelling older persons. *Journal of the American Geriatric Society, 51*(8), 1092-1098.
- Rejeski, W. J., Miller, M. E., Foy, C., Messier, S., & Rapp, S. (2001). Self-efficacy and the progression of functional limitations and self-reported disability in older adults with knee pain. *The Journal of Gerontology. Social Sciences, 56*(5), S261-S265.
- Reyes-Gibby, C. C., Aday, L., & Cleeland, C. (2002). Impact of pain on self-rated health in the community-dwelling older adults. *Pain, 95*(1-2), 75-82.
- Riley, J. L., & Gilbert, G. H. (2001). Orofacial pain symptoms: An interaction between age and sex. *Pain, 90*(3), 245-256.
- Riley, J. L., Gilbert, G. H., & Heft, M. W. (1999). Health care utilization by older adults in response to painful orofacial symptoms. *Pain, 81*(1-2), 67-75.

- Riley, J. L., Gilbert, G. H., & Heft, M. W. (2002a). Orofacial pain-related communication patterns: Sex and residential setting differences among community-dwelling adults. *Pain*, 99(3), 415-422.
- Riley, J. L., Gilbert, G. H., & Heft, M. W. (2002b). Orofacial pain: Racial and sex differences among older adults. *Journal of Public Health Dentistry*, 62(3), 132-139.
- Riley, J. L., Gilbert, G. H., & Heft, M. W. (2002c). Race/ethnic differences in health care use for orofacial pain among older adults. *Pain*, 100(1-2), 119-130.
- Riley, J. L., Tomar, S. L., & Gilbert, G. H. (2004). Smoking and smokeless tobacco: Increased risk for oral pain. *Journal of Pain*, 5(4), 218-225.
- Riley, J. L., Wade, J. B., Robinson, M. E., & Price, D. D. (2000). The stages of pain processing across the adult lifespan. *The Journal of Pain*, 1(2), 162-170.
- Roberto, K. A., & Reynolds, S. G. (2001). The meaning of osteoporosis in the lives of rural older women. *Health Care for Women International*, 22(6), 599-611.
- Roberto, K. A., & Reynolds, S. G. (2002). Older women's experiences with chronic pain: Daily challenges and self-care practices. *Journal of Women & Aging*, 14(3-4), 5-23.
- Roberts, R. O., Jacobson, D. J., Girman, C. J., Rhodes, T., Lieber, M. M., & Jacobsen, S. J. (2004). Low agreement between previous physician diagnosed prostatitis and national institutes of health chronic prostatitis symptom index pain measures. *The Journal of Urology*, 171(1), 279-283.
- Ross, M. M., Carswell, A., Hing, M., Hollingworth, G., & Dalziel, W. B. (2001). Seniors' decision making about pain management. *Journal of Advanced Nursing*, 35(3), 442-451.
- Rucker-Whitaker, C., Greenland, P., Liu, K., Chan, C., Guralnik, J. M., Criqui, M. H., Taylor, L., Pearce, W. H., & McDermott, M. (2004). Peripheral arterial disease in African Americans: Clinical characteristics, leg symptoms, and lower extremity functioning. *Journal of the American Geriatric Society*, 52(6), 922-930.
- Sarkisian, C. A., Hays, R. D., Berry, S. H., & Mangione, C. M. (2001). Expectations regarding aging among older adults and physicians who care for older adults. *Medical Care*, 39(9), 1025-1036.
- Schiff, M., & Minic, M. (2004). Comparison of the analgesic efficacy and safety of nonprescription doses of naproxen sodium and Ibuprofen in the treatment of osteoarthritis of the knee. *Journal of Rheumatology*, 31(7), 1373-1383.
- Scudds, R. J., & Ostbye, T. (2001). Pain and pain-related interference with function in older Canadians: The Canadian Study of Health and Aging. *Disability and Rehabilitation*, 23(15), 654-664.
- Scudds, R. J., & Robertson, J. M. (2000). Pain factors associated with physical disability in a sample of community-dwelling senior citizens. *The Journal of Gerontology. Medical Sciences*, 55(7), M393-M399.

- Sherman, A. M. (2003). Social relations and depressive symptoms in older adults with knee osteoarthritis. *Social Science & Medicine*, 56(2), 247-257.
- Sherman, K. J., Cherkin, D. C., Connelly, M. T., Erro, J., Savetsky, J. B., Davis, R. B., & Eisenberg, D. M., (2004). Complementary and alternative medical therapies for chronic low back pain: What treatments are patients willing to try? *BMC Complementary and Alternative Medicine*, 4(1), 9.
- Shmueli, Y., Baumgarten, M., Rovner, B., & Berlin, J. (2001). Predictors of improvement in health-related quality of life among elderly patients with depression. *International Psychogeriatrics*, 13(1), 63-73.
- Shrader, M. W., Draganich, L. F., Pottenger, L. A., & Piotrowski, G. A. (2004). Effects of knee pain relief in osteoarthritis on gait and stair-stepping. *Clinical Orthopaedics* (421), 188-193.
- Simmons, S. F., Ferrell, B. A., & Schnelle, J. F. (2002). Effects of a controlled exercise trial on pain in nursing home residents. *The Clinical Journal of Pain*, 18(6), 380-385.
- Stevens-Ratchford, R., & Lookingbill, J. (2004). Living well with arthritis: A phenomenological study of leisure occupations. *Activities, Adaptation & Aging*, 28(2), 35-55.
- Summaries for patients. Musculoskeletal pain and the progression of disability among older women with disability. (2001). *Annals of Internal Medicine*, 135(12), S-64.
- Tannenbaum, H., Berenbaum, F., Reginster, J. Y., Zacher, J., Robinson, J., Poor, G., Bliddal, H., Uebelhart, D., Adami, S., Navarro, F., Lee, A., Moore, A., & Gimona, A. (2004). Lumiracoxib is effective in the treatment of osteoarthritis of the knee: A 13 week, randomised, double blind study versus placebo and celecoxib. *Annals of the Rheumatic Diseases*, 63(11), 1419-1426.
- Tarzian, A. J., & Hoffmann, D. E. (2004). Barriers to managing pain in the nursing home: Findings from a statewide survey. *Journal of the American Medical Directors Association*, 5(2), 82-88.
- Taylor, L. J., & Herr, K. (2003). Pain intensity assessment: A comparison of selected pain intensity scales for use in cognitively intact and cognitively impaired African American older adults. *Pain Management Nursing*, 4(2), 87-95.
- Teno, J. M., Kabumoto, G., Wetle, T., Roy, J., & Mor, V. (2004). Daily pain that was excruciating at some time in the previous week: Prevalence, characteristics, and outcomes in nursing home residents. *Journal of the American Geriatrics Society*, 52(5), 762-767.
- Tran, F., Boggie, D. T., Delattre, M. L., Schaefer, M. G., Morreale, A. P., & Plowman, B. K. (2004). Therapeutic interchange involving replacement of rofecoxib or celecoxib with valdecoxib. *American Journal of Health-System Pharmacy*, 61(13), 1391-1394.
- Tsai, P. F., Tak, S., Moore, C., & Palencia, I. (2003). Testing a theory of chronic pain. *Journal of Advanced Nursing*, 43(2), 158-169.

- Unutzer, J., Ferrell, B., Lin, E. H., & Marmon, T. (2004). Pharmacotherapy of pain in depressed older adults. *Journal of the American Geriatric Society*, 52(11), 1916-1922.
- Villanueva, M. R., Smith, T. L., Erickson, J. S., Lee, A. C., & Singer, C. M. (2003). Pain Assessment for the Dementing Elderly (PADE): Reliability and validity of a new measure. *Journal of the American Medical Directors Association*, 4(1), 1-8.
- Vogt, M. T., Lauerma, W. C., Chirumbole, M., & Kuller, L. H. (2002). A community-based study of postmenopausal white women with back and leg pain: Health status and limitations in physical activity. *The Journal of Gerontology. Medical Sciences*, 57(8), M544-M550.
- Vogt, M. T., Simonsick, E. M., Harris, T. B., Nevitt, M. C., Kang, J. D., Rubin, S. M., Kritchevsky, S. B., & Newman, A. B., (2003). Neck and shoulder pain in 70- to 79-year-old men and women: Findings from the Health, Aging and Body Composition Study. *The Spine Journal*, 3(6), 435-441.
- Weiner, D., Peterson, B., & Keefe, F. (1999). Chronic pain-associated behaviors in the nursing home: Resident versus caregiver perceptions. *Pain*, 80(3), 577-588.
- Weiner, D., Peterson, B., Ladd, K., McConnell, E., & Keefe, F. (1999). Pain in nursing home residents: An exploration of prevalence, staff perspectives, and practical aspects of measurement. *The Clinical Journal of Pain*, 15(2), 92-101.
- Weiner, D. K., Haggerty, C. L., Kritchevsky, S. B., Harris, T., Simonsick, E. M., Nevitt, M., & Newman, A., (2003). How does low back pain impact physical function in independent, well-functioning older adults? Evidence from the Health ABC Cohort and implications for the future. *Pain Medicine*, 4(4), 311-320.
- Weiner, D. K., & Rudy, T. E. (2002). Attitudinal barriers to effective treatment of persistent pain in nursing home residents. *Journal of the American Geriatrics Society*, 50(12), 2035-2040.
- Weiner, D. K., Rudy, T. E., & Gaur, S. (2001). Are all older adults with persistent pain created equal? Preliminary evidence for a multiaxial taxonomy. *Pain Research & Management : The Journal of the Canadian Pain Society* , 6(3), 133-141.
- Weiner, D. K., Rudy, T. E., Glick, R. M., Boston, J. R., Lieber, S. J., Morrow, L. A., & Taylor, S. (2003). Efficacy of percutaneous electrical nerve stimulation for the treatment of chronic low back pain in older adults. *Journal of the American Geriatrics Society*, 51(5), 599-608.
- Weintraub, M. I., & Cole, S. P. (2004). Pulsed magnetic field therapy in refractory neuropathic pain secondary to peripheral neuropathy: Electrodiagnostic parameters--pilot study. *Neurorehabilitation and Neural Repair*, 18(1), 42-46.
- Wilcox, S., Brenes, G. A., Levine, D., Seveck, M. A., Shumaker, S. A., & Craven, T. (2000). Factors related to sleep disturbance in older adults experiencing knee pain or knee pain with radiographic evidence of knee osteoarthritis. *Journal of the American Geriatric Society*, 48(10), 1241-1251.

- Williams, D. A., Farrell, M. J., Cunningham, J., Gracely, R. H., Ambrose, K., Cupps, T., Mohan, N., & Clauw, D. J. (2004). Knee pain and radiographic osteoarthritis interact in the prediction of levels of self-reported disability. *Arthritis and Rheumatism*, 51(4), 558-561.
- Wolsko, P. M., Eisenberg, D. M., Simon, L. S., Davis, R. B., Walleczek, J., Mayo-Smith, M., Kaptchuk, T. J., & Phillips, R. S. (2004). Double-blind placebo-controlled trial of static magnets for the treatment of osteoarthritis of the knee: Results of a pilot study. *Alternative Therapies in Health and Medicine*, 10(2), 36-43.
- Won, A., Lapane, K., Gambassi, G., Bernabei, R., Mor, V., & Lipsitz, L. A. (1999). Correlates and management of nonmalignant pain in the nursing home. SAGE Study Group. Systematic Assessment of Geriatric drug use via Epidemiology. *Journal of the American Geriatrics Society*, 47(8), 936-942.
- Won, A. B., Lapane, K. L., Vallow, S., Schein, J., Morris, J. N., & Lipsitz, L. A. (2004). Persistent nonmalignant pain and analgesic prescribing patterns in elderly nursing home residents. *Journal of the American Geriatrics Society*, 52(6), 867-874.
- Wu, N., Miller, S. C., Lapane, K., & Gozalo, P. (2003). The problem of assessment bias when measuring the hospice effect on nursing home residents' pain. *Journal of Pain and Symptom Management*, 26(5), 998-1009.
- Wynne, C. F., Ling, S. M., & Remsburg, R. (2000). Comparison of pain assessment instruments in cognitively intact and cognitively impaired nursing home residents. *Geriatric Nursing*, 21(1), 20-23.
- Yamakawa, K., Tsai, C. K., Haig, A. J., Miner, J. A., & Harris, M. J. (2004). Relationship between ambulation and obesity in older persons with and without low back pain. *International Journal of Obesity and Related Metabolic Disorders*, 28(1), 137-143.
- Yan, L. L., Daviglus, M. L., Liu, K., Pirzada, A., Garside, D. B., Schiffer, L., Dyer, A. R., & Greenland, P. (2004). BMI and health-related quality of life in adults 65 years and older. *Obesity Research*, 12(1), 69-76.